

Identification of G-quadruplex-forming Sequences in Nucleocapsid Gene of SARS-CoV-2 Variants of Concern: An *in silico* Analysis

Saeedeh Ghazaey Zidanloo

Department of Cell and Molecular Biology, Faculty of Science, Kosar University of Bojnord, Bojnord, Iran

ARTICLE INFO

Article history:

Received 05 August 2023

Accepted 18 September 2023

Available online 01 October 2023

Keywords:

G-quadruplex

Nucleocapsid gene

SARS-CoV2

Variants of concern

Supplementary information:

Supplementary information for

this article is available at

<http://sc.journals.umz.ac.ir/>

*Corresponding authors:

✉ S. Ghazaey Zidanloo
Ghazaey@s@kub.ac.ir

p-ISSN 2423-4257

e-ISSN 2588-2589

ABSTRACT

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), as an enveloped RNA virus, has resulted in a global health threat. Recent studies emphasized that G-quadruplex structures are intrinsic obstacles to genome replication and targeting them in viral genomes could be a novel antiviral strategy to develop antiviral agents. The genomic RNA of SARS-CoV-2 codes for 29 proteins. One of them is the nucleocapsid protein with multiple functions, which is crucial for several steps of the viral life cycle. Here, we have analyzed putative G-quadruplex sequences (PQSs) in the Nucleocapsid gene of SARS-CoV2 and its variants of concern using a bioinformatics tool. The results showed that the number, position, and G-scores of PQSs were similar in Wuhan-hu-1 and Alpha, Beta, and Gamma variants. The main difference was observed in the Delta variant, in which a PQS was deleted at position 630 of the gene, which is the top-ranked highly conserved G-quadruplex. The Omicron variant had this PQS back at position 621 as it had acquired several mutations. In addition, there is also a unique R203M mutation, in the N protein of the Delta variant that leads to increased RNA packaging, replication, and severe COVID-19. We proposed that the R203M mutation has led to G>T substitution and loss of the top-ranked highly conserved PQS in the N gene of the Delta variant. Therefore, due to the loss of this important PQS or indeed an obstacle to viral replication, the Delta variant could exhibit higher reproduction and pathogenicity than other variants of concern.

© 2024 University of Mazandaran

Please cite this paper as: Ghazaey Zidanloo, S. (2024). Identification of G-quadruplex-forming sequences in nucleocapsid gene of SARS-CoV-2 variants of concern: an *in silico* analysis. *Journal of Genetic Resources*, 10(1), doi: [10.22080/jgr.2024.25850.1370](https://doi.org/10.22080/jgr.2024.25850.1370)

Introduction

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has spread across the world rapidly since the end of 2019 and has resulted in an epidemic of viral pneumonia, and a global health threat. Although the global scientific community has not found any effective antiviral treatment, researchers assist in creating novel COVID-19 diagnostic and treatment methods. SARS-CoV-2 as an enveloped RNA virus, along with other phylogenetically related species could infect huge numbers of vertebrates (Astuti, 2020; Díaz, 2020). Up to now, five SARS-CoV-2 variants of concern (VOCs) have been reported including B.1.1.7 (Alpha), B.1.351

(Beta), P.1 (Gamma), B.1.617.2 (Delta) and B.1.1.529 (Omicron). Emerging evidence revealed increased transmissibility and virulence as well as, reduced neutralization, which would be even more dangerous and troublesome. It should be noted that the SARS-CoV-2 genome undergoes frequent and quick mutation during its life which might serve as potential therapeutic targets against different SARS-CoV-2 variants. Each VOC is defined by a combination of point mutations across the SARS-CoV-2 genome. Although some mutations are common among different VOCs, there are some novel mutations observed in one individual variant (Telenti *et al.*, 2022). Delta, as one of the most concerned



variants, is related to a greater risk of hospitalization, which shared some mutations with other variants, and owned its special mutations, which might explain the increase in transmission and virulence (Zhan *et al.*, 2022). Under these circumstances, a better understanding of the VOC's genetic variations and their clinical implications is necessary to provide new aspects for antiviral treatments.

The SARS-CoV-2 genomic RNA is ~30 kb long encoding about 25 non-structural proteins (nsps) and accessory proteins and also, four structural proteins including spike (S), envelope (E), membrane (M), and nucleocapsid (N) (Chan *et al.*, 2020; Huston *et al.*, 2021). Exclusively, one of the most conserved and stable structural proteins is the N protein which is composed of 419 amino acids and is encoded by the ninth ORF of the virus (He *et al.*, 2004). Nucleocapsid protein has multiple functions, participates in helical ribonucleoprotein formation in the packaging of the genomic RNA, regulation of RNA synthesis during viral replication and transcription, and control of metabolism in infected cells. Therefore, this protein is crucial for various steps of the viral life cycle (Chang *et al.*, 2014; Narayanan *et al.*, 2003).

Recent studies emphasized that secondary structural elements such as G-quadruplexes, as relatively highly conserved structures, could be used as potential antiviral targets (Zhai *et al.*, 2022b). RNA G-quadruplexes (RG4) that can be formed in guanine (G) rich in RNAs, contain two or more layers of G-quartets constructed in a planar arrangement by Hoogsteen hydrogen bonding (Agarwala *et al.*, 2015; Varshney *et al.*, 2020). It can be suggested that RG4 might fold and stabilize the RNA motif appropriately to prevent viral replication, transcription, and translation (Abiri *et al.*, 2021; Lyu *et al.*, 2021; Metifiot *et al.*, 2014; Ruggiero *et al.*, 2021; Xu *et al.*, 2021). There is increasing evidence that G-quadruplex structures play a regulatory role in gene transcription in humans and other organisms and are important potential therapeutic targets (Rhodes and Lipps, 2015; Zidanloo *et al.*, 2016)

This study, for the first time, analyzed the putative G-quadruplexes-forming sequences (PQS) *in silico* in nucleocapsid gene of SARS-CoV-2 different VOCs and the PQSs possible

impact on the packaging of the viral genome and viral life cycle. Therefore, it can be suggested that the G-quadruplex motifs presenting in the N gene, might play a fundamental role in regulating its function and serving as potential targets for SARS-CoV-2 treatment.

Methods

SARS-CoV-2 genomic sequences

The genomic sequence of the SARS-CoV-2 Wuhan Hu-1 RNA reference genome (MN908947) and other global variants of concern were downloaded from EpiCoV database of Global Initiative on Sharing All Influenza Data (GISAID) (<https://www.gisaid.org/>) using automatic search function feeding information for geographical location, SARS-CoV-2 lineage and sample collection and sequence reporting dates (up to 2023-19-07). Afterward, the N gene sequence of different strains was extracted for further investigation.

G-quadruplex mapping

G-quadruplex sequences of the N gene are mapped on the SARS-CoV2 genome via the website QGRS mapper software (<https://bioinformatics.ramapo.edu/QGRS/analyze.php>) (Kikin *et al.*, 2006). PQS of a maximum length of 30 bases and loop consisting of any length from 0 to 30 base sequences are considered. As an output, this software gives the list of detected PQSs (unique and overlapping sequences) with the scoring data. The possibility of the G-quadruplex structure formation for each sequence was gained as G-scores. In addition, the position of each PQS sequence in the forward and reverse strands of the SARS-CoV2 genome was listed as software output.

Result

Identification of RNA G-quadruplexes in SARS-CoV-2 N gene.

To explore the potential involvement of RNA G-quadruplexes in SARS-CoV-2 VOCs, an *in silico* method with online software was used (Kikin *et al.*, 2006) which offered a G-score for each PQS in the virus genome. The PQSs were predicted in the SARS-CoV-2 genome by the QGRS mapper algorithm. In the 29.9 kb genome

of SARS-CoV-2, the PQSs in the Nucleocapsid gene (nt. 28274-29533) of VOCs were analyzed. Consequently, any position with a higher probability of forming RNA G-quadruplex

structures had a higher score so, the presence of potential sequence stretch that could form a G-quadruplex structure was listed in Table 1 (Supplement 1-6).

Table 1. Putative G-quadruplex sequences in Nucleocapsid gene of SARS-CoV-2 variants of concern

WHO nomenclature	Lineage	Date of designation	Emergence	QGRS/PQS position ^a (G-Score) [Overlaps not included]	QGRS/PQS position (G-Score) [Overlaps included]
Wuhan-hu-1	Reference	Dec, 2019	China	508 (9), 630 (18) , 850 (14), 961 (11)	508 (9), 508 (8), 608 (3), 630 (18) , 850 (14), 859 (7), 961 (11), 961 (10)
Alpha (α)	B.1.1.7	Dec, 2020	UK	508 (9), 630 (18) , 850 (14), 961 (11)	508 (9), 508 (8), 630 (18) , 850 (14), 859 (7), 961 (11), 961 (10)
Beta (β)	B.1.351	Dec, 2020	South Africa	508 (9), 630 (18) , 850 (14), 961 (11)	508 (9), 508 (8), 608 (3), 630 (18) , 850 (14), 859 (7), 961 (11), 961 (10)
Gamma (P.1)	B.1.1.128	Jan, 2021	Brazil	508 (9), 630 (18) , 850 (14), 961 (11)	508 (9), 508 (8), 630 (18) , 850 (14), 859 (7), 961 (11), 961 (10)
Delta (δ)	B.1.617.2	May, 2021	India	178 (7), 508 (9), 850 (14), 961 (11)	178 (7), 183 (7), 508 (9), 508 (8), 850 (14), 859 (7), 961 (11), 961 (10)
Omicron (o)	B.1.1.529	Nov, 2021	South Africa	73 (8), 499 (9), 621 (18) , 841 (14), 952 (11)	73 (8), 88 (3), 499 (9), 499 (8), 621 (18) , 841 (14), 850 (7), 952 (11), 952 (10)

a: nucleotide position

We found several PQSs in the N gene of the SARS-CoV2 and the VOCs with the acceptable potential to form a G-quadruplex structure. Even though PQSs were confirmed to be present near the transcription start site of many genes (Kikin *et al.*, 2006; Perrone *et al.*, 2017), the PQSs of the N gene were not located in these regions. Without considering overlapping sequences, there are four PQS in the N gene of VOCs except for the Omicron with five PQSs regions. If overlapping sequences are considered, up to nine PQS regions for SARS-CoV-2 VOCs are observed (Supplementary Material). These regions all fit the G2N1-7G2N1-7G2N1-7G2 formula, indicating the potential to adopt the non-canonical and metastable RG4 structures with 2-quartet (Ji *et al.*, 2021; Zhao *et al.*, 2021). This analysis revealed that in all PQSs of the N gene, there are regions with four contiguous GG runs and lengths of 15-30 nucleotides (See Supplementary Supplement 1-6; also Table 1). The QGRS mapper results revealed that the number, position, and G-scores of PQSs were similar in Wuhan-hu-1 SARS-CoV-2 and the three VOCs Alpha, Beta, and Gamma. The differences were observed in the Delta variant in which a PQS at position 630 (RG-1) was deleted (Supplement 5). The Omicron variant exhibited this PQS again at position 621 due to the several mutations it acquired. This PQS showed the highest G-score (G-score =18) in both analyses considering overlap and not considering overlap (bold in Table 1, Supplement 6).

Discussion

SARS-CoV-2 has been circulating in the human population for three years and has infected hundreds of millions of people. Therefore, understanding the molecular mechanism that could affect its gene expression in host cells is a prerequisite for developing a successful and effective treatment for COVID-19.

This is the first study to analyze the presence of PQSs in different VOCs of SARS-CoV-2. Our result confirmed that the highest ranked PQS (RG-1, G-score = 18) was present at position 630 (621 for Omicron) in all VOCs except the Delta variant (Table 1). Recent studies identified G-quadruplex motifs in the SARS-CoV-2 genome and also found that some PQSs were well-conserved in a wide range of coronaviruses (Cui and Zhang, 2020; Panera *et al.*, 2020; Ji *et al.*, 2021). Also, Belmonte-Reche *et al.* identified PQSs with scores higher than 20, and conserved PQSs in SARS-CoV-2, SARS-CoV, and Bat-CoV (Belmonte-Reche *et al.*, 2021). In another study, Zhang *et al.* obtained 24 PQSs with three prediction tools nine of which were scattered on the negative strand (Zhang *et al.*, 2020). According to the importance of G-quadruplex structures in promoter regions (Zidanloo *et al.*, 2016), Maiti focused on those SARS-CoV-2 genes that have PQS sequences upstream at the putative promoter region and identified PQSs upstream of at least 16 genes of the virus (Maiti, 2022). It is noteworthy that the expression of

these genes might be controlled via genome rearrangements or stability and by preventing RNA splicing or translation (Machida *et al.*, 2020). Although there are no such potential sequences upstream of the SARS-CoV-2 N gene, it is notable that the PQSs in any position of a gene can modify gene expression. Besides, the results of a recent study showed that the higher the number of PQS, the more the replication rate of the viruses is affected. Thus, the fact that SARS-CoV-2 contains fewer PQSs than SARS-CoV seems to be consistent with the fact that SARS-CoV-2 has a stronger replication capacity and faster spread (Zhai *et al.*, 2022a).

Lately, seven conserved RG4 sequences have been reported in the genomes of SARS-CoV and SARS-CoV-2, three in the S coding region, two in the nsP1 coding region, and the other two in the nsP10 and N coding regions (Cui and Zhang, 2020). In addition, another study analyzed more than 200,000 SARS-CoV-2 genome sequences from five continents and found PQSs at six positions with a high degree of conservation of more than 90%. Interestingly one of them was a PQS at N protein-coding sequence (Carvalho *et al.*, 2021) which is the top-ranked PQS of the N gene (RG-1) (Table 1). It has been confirmed that RG-1 can form a stable parallel G-quadruplex conformation with some ligands can interact and potentially over-stabilize it to significantly reduce protein levels of SARS-CoV-2 N protein by inhibiting viral genome expression both *in vitro* and *in vivo* and impairing the viral life cycle (Micolot *et al.*, 2021; Mukherjee *et al.*, 2022; Zhao *et al.*, 2021).

Several mutations were observed in the N protein of SARS-CoV-2 VOCs, including D3L (aspartic acid to leucine substitution at position 3), R203K (arginine to lysine substitution at position 203), R203M (arginine to methionine substitution at position 203), G204R (glycine to arginine substitution at position 204), S194L (serine to leucine substitution at position 194) and S235F (serine to phenylalanine substitution at position 235), T205I, (threonine to isoleucine substitution at position 205). It is worth noting that R203M, G28881T substitution in the Delta genome, is a common mutation that occurred in the N protein and is significantly associated with the Delta variant (Ravi *et al.*, 2022; Khetrani and Mustafa, 2023). In another study, Li and his co-

workers pointed out that the Delta variant may have a faster viral replication rate and is more infectious in the early stage of infection (viral RNA concentration is increased by a factor of 1000 in patients) (Li *et al.*, 2022). In addition, several studies have found a link between the R203M mutation and severe COVID-19 (Alsuwairi *et al.*, 2023; Mourier *et al.*, 2022; Zhao *et al.*, 2022). Interestingly, Syed *et al.*, showed that the R203M mutation in the N gene of Delta variants leads to increased RNA packaging and replication (more than 50-fold) (Syed *et al.*, 2021).

Conclusions

The RNA G-quadruplexes block viral replication in SARS-CoV-2 and could inhibit viral proliferation in SARS-CoV-2 VOCs (Fig. 1). We propose that the unique R203M mutation in the Delta variant could result in the loss of the highly conserved PQS (RG-1) of the N gene. The mutation R203M (G>T) is located at position 28881 of the Delta variant genome and led to the loss of one of the guanines in this region, which is involved in the formation of the G-quadruplex structure (Supplement 7). Therefore, the Delta variant may have effective and efficient replication and higher pathogenicity than other variants of concern due to the absence of this important PQS or an obstacle to viral replication (Fig. 2). Since the presence of this conserved PQS could play a regulatory role in the viral genome, the highest-ranking PQS in the N gene (RG-1) is an interesting topic for further research and could be the most attractive candidate for further investigation of SARS-CoV-2 gene expression. Exploring such roles by designing future experiments with PQS could be extremely useful in understanding whether these sequences modulate the expression of SARS-CoV2 genes and support viral amplification for their survival in human cells. Nevertheless, G-quadruplex nucleic acid structures are relatively difficult to study *in vivo*, and only a limited number of structures have been solved to date (Ji *et al.*, 2021).

Conflicts of interest

There was no conflict of interest in the report.




Fig. 1. RNA G-quadruplex could inhibit viral proliferation in SARS-CoV-2 VOCs: A) Schematic representation of the SARS-CoV-2 genome and the N gene. B) The RG-1 sequence in the N gene, the structure of its G-quartet, and its parallel G-quadruplex at position 630. C) Stalling the polymerase via the formation of a top-ranked G-quadruplex structure (RG-1) could reduce viral replication.




Fig. 2. R203M takes place in position 203 of the N protein in the Delta variant, which is nucleotide 28881 of the virus genome. It exactly overlaps with the formation of the top-ranked PQS. Guanine-rich sequences with four guanine repeats tend to fold intramolecularly into DNA quadruplexes. This guanine at 28881, plays an important role in the formation of the PQS, and the conversion of G to T reduces the possibility of the formation of a G-quadruplex structure.

Funding

This work was supported by the Kosar University of Bojnord under Grant number [NO.0102262653].

Conflicts of interest

Authors declare that there is no conflict of interest regarding the publication of this article.

References

- Abiri, A., Lavigne, M., Rezaei, M., Nikzad, S., Zare, P., Mergny, J. L., & Rahimi, H. R. (2021). Unlocking G-quadruplexes as antiviral targets. *Pharmacological Reviews*, 73(3), 897-923. <https://doi.org/10.1124/pharmrev.120.000230>
- Agarwala, P., Pandey, S., & Maiti, S. (2015). The tale of RNA G-quadruplex. *Organic & Biomolecular Chemistry*, 13(20), 5570-5585. <https://doi.org/10.1039/c4ob02681k>
- Alsuwairi, F. A., Alsaleh, A. N., Alsanea, M. S., Al-Qahtani, A. A., Obeid, D., Almaghrabi, R. S., ... & Alhamlan, F. S. (2023). Association of SARS-CoV-2 nucleocapsid protein mutations with patient demographic and clinical characteristics during the Delta and Omicron waves. *Microorganisms*, 11(5), 1288. <https://doi.org/10.3390/microorganisms11051288>
- Astuti, I. (2020). Severe Acute respiratory syndrome coronavirus 2 (SARS-CoV-2): an overview of viral structure and host response. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 14(4), 407-412. <https://doi.org/10.1016/j.dsx.2020.04.020>
- Belmonte-Reche, E., Serrano-Chacon, I., Gonzalez, C., Gallo, J., & Banobre-Lopez, M. (2021). Potential G-quadruplexes and i-motifs in the SARS-CoV-2. *PLoS One*, 16(6), e0250654. <https://doi.org/10.1371/journal.pone.0250654>
- Carvalho, J., Lopes-Nunes, J., Figueiredo, J., Santos, T., Miranda, A., Riscado, M., ... & Cruz, C. (2021). Molecular beacon assay development for severe acute respiratory syndrome coronavirus 2 detection. *Sensors*, 21(21), 7015. <https://doi.org/10.3390/s21217015>
- Chan, J. F. W., Kok, K. H., Zhu, Z., Chu, H., To, K. K. W., Yuan, S., & Yuen, K. Y. (2020). Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerging Microbes & Infections*, 9(1), 221-236. <https://doi.org/10.1080/22221751.2020.1719902>
- Chang, C. K., Hou, M. H., Chang, C. F., Hsiao, C. D., & Huang, T. H. (2014). The SARS coronavirus nucleocapsid protein—forms and functions. *Antiviral Research*, 103, 39-50. <https://doi.org/10.1016/j.antiviral.2013.12.009>
- Cui, H., & Zhang, L. (2020). G-quadruplexes are present in human coronaviruses including SARS-CoV-2. *Frontiers in Microbiology*, 11, 567317. <https://doi.org/10.3389/fmicb.2020.567317>
- Díaz, J. (2020). SARS-CoV-2 molecular network structure. *Frontiers in Physiology*, 11, 870. <https://doi.org/10.3389/fphys.2020.00870>
- He, Y., Zhou, Y., Wu, H., Kou, Z., Liu, S., & Jiang, S. (2004). Mapping of antigenic sites on the nucleocapsid protein of the severe acute respiratory syndrome coronavirus. *Journal of Clinical Microbiology*, 42(11), 5309-5314. <https://doi.org/10.1128/jcm.42.11.5309-5314.2004>
- Huston, N. C., Wan, H., Strine, M. S., Tavares, R. D. C. A., Wilen, C. B., & Pyle, A. M. (2021). Comprehensive *in vivo* secondary structure of the SARS-CoV-2 genome reveals novel regulatory motifs and mechanisms. *Molecular Cell*, 81(3), 584-598. <https://doi.org/10.1016/j.molcel.2020.12.041>
- Ji, D., Juhas, M., Tsang, C. M., Kwok, C. K., Li, Y., & Zhang, Y. (2021). Discovery of G-quadruplex-forming sequences in SARS-CoV-2. *Briefings in Bioinformatics*, 22(2), 1150-1160. <https://doi.org/10.1093/bib/bbaa114>
- Khetran, S. R., & Mustafa, R. (2023). Mutations of SARS-CoV-2 structural proteins in the alpha, beta, gamma, and delta variants: bioinformatics analysis. *JMIR Bioinformatics and Biotechnology*, 4(1), e43906. <https://doi.org/10.2196/43906>
- Kikin, O., D'Antonio, L., & Bagga, P. S. (2006). QGRS Mapper: a web-based server for predicting G-quadruplexes in nucleotide sequences. *Nucleic Acids Research*, 34(suppl_2), W676-W682. <https://doi.org/10.1093/nar/gkl253>
- Li, B., Deng, A., Li, K., Hu, Y., Li, Z., Shi, Y., ... & Lu, J. (2022). Viral infection and transmission in a large, well-traced outbreak

- caused by the SARS-CoV-2 Delta variant. *Nature Communications*, 13(1), 460. <https://doi.org/10.1038/s41467-022-28089-y>
- Lyu, K., Chow, E. Y. C., Mou, X., Chan, T. F., & Kwok, C. K. (2021). RNA G-quadruplexes (rG4s): genomics and biological functions. *Nucleic Acids Research*, 49(10), 5426-5450. <https://doi.org/10.1093/nar/gkab187>
- Machida, S., Depierre, D., Chen, H. C., Thenin-Houssier, S., Petitjean, G., Doyen, C. M., ... & Benkirane, M. (2020). Exploring histone loading on HIV DNA reveals a dynamic nucleosome positioning between unintegrated and integrated viral genomes. *Proceedings of the National Academy of Sciences*, 117(12), 6822-6830. <https://doi.org/10.1073/pnas.1913754117>
- Maiti, A. K. (2022). Identification of G-quadruplex DNA sequences in SARS-CoV2. *Immunogenetics*, 74(5), 455-463. <https://doi.org/10.1007/s00251-022-01257-6>
- Métifiot, M., Amrane, S., Litvak, S., & Andreola, M. L. (2014). G-quadruplexes in viruses: function and potential therapeutic applications. *Nucleic Acids Research*, 42(20), 12352-12366. <https://doi.org/10.1093/nar/gku999>
- Miclot, T., Hognon, C., Bignon, E., Terenzi, A., Marazzi, M., Barone, G., & Monari, A. (2021). Structure and dynamics of RNA guanine quadruplexes in SARS-CoV-2 genome. original strategies against emerging viruses. *The Journal of Physical Chemistry Letters*, 12(42), 10277-10283. <https://doi.org/10.1021/acs.jpcclett.1c03071>
- Mourier, T., Shuaib, M., Hala, S., Mfarrej, S., Alofi, F., Naeem, R., ... & Pain, A. (2022). SARS-CoV-2 genomes from Saudi Arabia implicate nucleocapsid mutations in host response and increased viral load. *Nature Communications*, 13(1), 601. <https://doi.org/10.1038/s41467-022-28287-8>
- Mukherjee, S. K., Knop, J. M., & Winter, R. (2022). Modulation of the conformational space of SARS-CoV-2 RNA quadruplex RG-1 by cellular components and the amyloidogenic peptides α -Synuclein and hIAPP. *Chemistry-A European Journal*, 28(9), e202104182. <https://doi.org/10.1002/chem.202104182>
- Narayanan, K., Kim, K. H., & Makino, S. (2003). Characterization of N protein self-association in coronavirus ribonucleoprotein complexes. *Virus Research*, 98(2), 131-140. <https://doi.org/10.1016/j.virusres.2003.08.021>
- Panera, N., Tozzi, A. E., & Alisi, A. (2020). The G-quadruplex/helicase world as a potential antiviral approach against COVID-19. *Drugs*, 80(10), 941-946. <https://doi.org/10.1007/s40265-020-01321-z>
- Perrone, R., Lavezzo, E., Riello, E., Manganelli, R., Palù, G., Toppo, S., ... & Richter, S. N. (2017). Mapping and characterization of G-quadruplexes in *Mycobacterium tuberculosis* gene promoter regions. *Scientific Reports*, 7(1), 5743. <https://doi.org/10.1038/s41598-017-05867-z>
- Ravi, V., Swaminathan, A., Yadav, S., Arya, H., & Pandey, R. (2022). SARS-CoV-2 variants of concern and variations within their genome architecture: does nucleotide distribution and mutation rate alter the functionality and evolution of the virus? *Viruses*, 14(11), 2499. <https://doi.org/10.3390/v14112499>
- Rhodes, D., & Lipps, H. J. (2015). G-quadruplexes and their regulatory roles in biology. *Nucleic Acids Research*, 43(18), 8627-8637. <https://doi.org/10.1093/nar/gkv862>
- Ruggiero, E., Zanin, I., Terreri, M., & Richter, S. N. (2021). G-quadruplex targeting in the fight against viruses: an update. *International Journal of Molecular Sciences*, 22(20), 10984. <https://doi.org/10.3390%2Fijms222010984>
- Syed, A. M., Taha, T. Y., Tabata, T., Chen, I. P., Ciling, A., Khalid, M. M., ... & Doudna, J. A. (2021). Rapid assessment of SARS-CoV-2-evolved variants using virus-like particles. *Science*, 374(6575), 1626-1632. <https://doi.org/10.1126/science.abl6184>
- Telenti, A., Hodcroft, E. B., & Robertson, D. L. (2022). The evolution and biology of SARS-CoV-2 variants. *Cold Spring Harbor Perspectives in Medicine*, 12(5), a041390. <https://doi.org/10.1101/cshperspect.a041390>
- Varshney, D., Spiegel, J., Zyner, K., Tannahill, D., & Balasubramanian, S. (2020). The regulation and functions of DNA and RNA G-quadruplexes. *Nature Reviews Molecular Cell Biology*, 21(8), 459-474. <https://doi.org/10.1038/s41580-020-0236-x>
- Xu, J., Huang, H., & Zhou, X. (2021). G-quadruplexes in neurobiology and virology: functional roles and potential therapeutic approaches. *Journal of the American*

- Chemical Society Au*, 1(12), 2146-2161. <https://doi.org/10.1021/jacsau.1c00451>
- Zhai, L. Y., Liu, J. F., Zhao, J. J., Su, A. M., Xi, X. G., & Hou, X. M. (2022a). Targeting the RNA G-Quadruplex and protein interactome for antiviral therapy. *Journal of Medicinal Chemistry*, 65(15), 10161-10182. <https://doi.org/10.1021/acs.jmedchem.2c00649>
- Zhai, L. Y., Su, A. M., Liu, J. F., Zhao, J. J., Xi, X. G., & Hou, X. M. (2022b). Recent advances in applying G-quadruplex for SARS-CoV-2 targeting and diagnosis: A review. *International Journal of Biological Macromolecules*. <https://doi.org/10.1016/j.ijbiomac.2022.09.152>
- Zhan, Y., Yin, H., & Yin, J. Y. (2022). B. 1.617. 2 (Delta) Variant of SARS-CoV-2: features, transmission and potential strategies. *International Journal of Biological Sciences*, 18(5), 1844. <https://doi.org/10.7150/ijbs.6-6881>
- Zhang, R., Xiao, K., Gu, Y., Liu, H., & Sun, X. (2020). Whole genome identification of potential G-quadruplexes and analysis of the G-quadruplex binding domain for SARS-CoV-2. *Frontiers in Genetics*, 11, 587829. <https://doi.org/10.3389/fgene.2020.587829>
- Zhao, C., Qin, G., Niu, J., Wang, Z., Wang, C., Ren, J., & Qu, X. (2021). Targeting RNA G-quadruplex in SARS-CoV-2: a promising therapeutic target for COVID-19? *Angewandte Chemie*, 133(1), 436-442. <https://doi.org/10.1002/anie.202011419>
- Zhao, L. P., Roychoudhury, P., Gilbert, P., Schiffer, J., Lybrand, T. P., Payne, T. H., ... & Geraghty, D. E. (2022). Mutations in viral nucleocapsid protein and endoRNase are discovered to be associated with COVID19 hospitalization risk. *Scientific Reports*, 12(1), 1206. <https://doi.org/10.1038/s41598-021-04376-4>
- Zidanloo, S. G., Hosseinzadeh Colagar, A., Ayatollahi, H., & Raoof, J. B. (2016). Downregulation of the WT 1 gene expression via TMPyP4 stabilization of promoter G-quadruplexes in leukemia cells. *Tumor Biology*, 37, 9967-9977. <https://doi.org/10.1007/s13277-016-4881-9>